

Bigger Shrimp

Optimizing Strategies For Growing Larger *L. vannamei*

Shaun M. Moss, Ph.D
smoss@oceanicinstitute.org

Clete A. Otoshi, M.S.

The Oceanic Institute
41-202 Kalaniana'ole Highway
Waimanola, Hawaii 96795 USA

PingSun Leung, Ph.D.
Department of Molecular Biosciences
And Bioengineering
University of Hawaii at Manoa
Honolulu, Hawaii USA

Although the giant tiger prawn, *Penaeus monodon*, is the most common shrimp species cultured in Asia, its dominance has faded in recent years as many Asian farmers stocked their ponds with Pacific white shrimp, *Litopenaeus vannamei*. Now more *L. vannamei* are produced in Asia than the Western Hemisphere, contributing to a glut of white shrimp and significant drops in shrimp prices.

Alternative Production Strategies

Shrimp farmers in the United States and elsewhere are considering alternative production and harvesting strategies to avoid direct competition with foreign imports from Asia, including the production of larger white shrimp. However, there is little published information on growth curves and feeding rates for larger *L. vannamei*, and even less information on optimal harvesting strategies.

O.I. Research

In an attempt to grow large shrimp at high densities under biosecure conditions, researchers at the Oceanic Institute stocked specific pathogen-free juvenile *L. vannamei* in a 58.4-m² raceway (Figure 1) at a density of 705 shrimp/m³. After 86 days, shrimp survival in the phase 1 study was 70.3%, mean harvest weight was 17.9 g, and mean growth rate was 1.37 g/week (Table 1). Production was 8.9

The Oceanic Institute has grown large white shrimp at high density in a biosecure raceway.



Summary:

Pressure from Asia is leading Western shrimp farmers to consider production and harvesting alternatives. Research at the Oceanic Institute found that white shrimp grown in a raceway at high density could reach 34-g sizes in 155 days with minimal water use. A two-phase harvesting approach proved more profitable than single harvests.

kg/m³, with mean water exchange throughout the trial of less than 2%/day.

Immediately following the first harvest, about 24% of the harvested shrimp were restocked into the raceway, which contained 55% of the original water, at 118 shrimp/m³. In this

phase 2 work, shrimp survival was 87.8% after 69 days, mean harvest weight was 34.2 g, and mean growth rate was 1.65 g/week.

During the second growout phase, shrimp exhibited sexual growth dimorphism. Females grew 1.88 g/week and were harvested at 36.9 g, where-




Figure 1. Shrimp growth over time in a biosecure raceway. Note the significant increase in growth immediately after the partial harvest at week 12.

Table 1. Performance of *L. vannamei* grown in a raceway. About 24% of the shrimp harvested from phase 1 were restocked for phase 2.

Parameter	Phase 1 Trial	Phase 2 Trial
Stocking density (shrimp/m ³)	705	118
Trial duration (days)	86	69
Stocking weight (g)	1.06	17.90
Harvest weight (g)	17.9	34.20
Survival (%)	70.3	87.80
Growth rate (g/week)	1.37	1.65
Production (kg/m ³)	8.9	3.50
Water use (l/kg shrimp)	219	229

as males grew 1.42 g/week and were harvested at 31.5 g. The trial indicated shrimp can be grown to 34.2 g in 155 days at high densities with minimal water use.

Compensatory Growth

Interestingly, weekly growth rates increased from 0.95 g during week 12, immediately prior to the first harvest, to 2.6 g during the first two weeks after the initial harvest (Figure 1). This dramatic increase suggested that *L. vannamei* may exhibit compensatory growth.

Compensatory growth occurs when an organism exhibits faster than normal growth during a period of recovery from growth restriction. In some fish, compensatory growth can be so intense that starved and refed individuals exhibit higher mean growth rates than those fed continuously.

In this case, growth restriction likely resulted from the high stocking density, as growth is density-dependent in penaeid shrimp. Earlier work at the Oceanic Institute showed that *L. vannamei* exhibit a rapid and significant increase in RNA concentration and RNA/DNA ratio following the reintroduction of feeding after several days of starvation. However, this was the first indication that the species may exhibit compensatory growth when relieved of density pressure.

Optimal Harvesting

From a farm management perspective, it may be possible to exploit this compensatory growth through partial harvesting strategies to increase production and profitability.

A partial budget analysis conducted by the Oceanic Institute using data from the growout trial indicated that partial harvesting was 8.8% more profitable than a single harvest if phase 2 shrimp were harvested



after four weeks. In contrast, if phase 2 were harvested after 10 weeks, there would have been a 5.9% loss in net revenue compared to a single harvest. In this case, the increase in yield for partial harvesting did not compensate for the opportunity loss of running more regular cycles.

The partial budget analysis used a unit of analysis of a single raceway and assumed 3.87 crops/year for the single-harvest scenario, 3.01 crops for the four-week phase 2 approach, and 2.25 crops with a phase 2 of 10 weeks. Prices for head-off shrimp with 2:1 tail/head ratios were based on prices from U.S. \$2.58/lb for 36/40 shrimp to \$6.50/lb for UN/15 shrimp. Feed cost was assumed to be \$0.70/kg, with other costs considered similar among the scenarios. Partial harvesting would increase harvesting costs somewhat.

Although this analysis suggested partial harvesting can be beneficial, the strategy has not been optimized. Growth curves and feeding rates for larger shrimp under different production scenarios are needed to develop efficient harvest models. Ultimately, these models can be used by farmers to increase production and improve profitability.